



# USE OF GLUCOSE-LOWERING MEDICATIONS IN THE MANAGEMENT OF TYPE 2 DIABETES

HEALTHY LIFESTYLE BEHAVIORS; DIABETES SELF-MANAGEMENT EDUCATION AND SUPPORT (DSMES); SOCIAL DETERMINANTS OF HEALTH (SDOH)



**Figure 9.3**—Use of glucose-lowering medications in the management of type 2 diabetes. ACEi, angiotensin-converting enzyme inhibitor; ACR, albumin-to-creatinine ratio; ARB, angiotensin receptor blocker; ASCVD, atherosclerotic cardiovascular disease; CGM, continuous glucose monitoring; CKD, chronic kidney disease; CV, cardiovascular; CVD, cardiovascular disease; CVOT, cardiovascular outcomes trial; DPP-4i, dipeptidyl peptidase 4 inhibitor; eGFR, estimated glomerular filtration rate; GLP-1 RA, glucagon-like peptide 1 receptor agonist; HF, heart failure; HFpEF, heart failure with preserved ejection fraction; HFrEF, heart failure with reduced ejection fraction; HFrEF, hospitalization for heart failure; MACE, major adverse cardiovascular events; MI, myocardial infarction; SDOH, social determinants of health; SGLT2i, sodium-glucose cotransporter 2 inhibitor; TZD, type 2 diabetes; TZD, thiazolidinedione. Adapted from Davies et al. (45).

Table 9.2—Medications for lowering glucose, summary of characteristics

	Efficacy <sup>1</sup>	Hypoglycemia	Weight change <sup>2</sup>	CV effects		Renal effects		Oral/SQ	Cost	Clinical considerations	
				Effect on MACE	HF	Progression of DKD	Dosing/use considerations*				
<b>Metformin</b>	High	No	Neutral (potential for modest loss)	Potential benefit	Neutral	Neutral	<ul style="list-style-type: none"> <li>Contraindicated with eGFR &lt;30 mL/min per 1.73 m<sup>2</sup></li> </ul>	Oral	Low	<ul style="list-style-type: none"> <li>GI side effects common; to mitigate GI side effects, consider slow dose titration, extended release formulations, and administration with food</li> <li>Potential for vitamin B12 deficiency; monitor at regular intervals</li> </ul>	
<b>SGLT2 inhibitors</b>	Intermediate to high	No	Loss (intermediate)	Benefit: canagliflozin, empagliflozin	Benefit: canagliflozin, dapagliflozin, empagliflozin, ertugliflozin	Benefit: canagliflozin, dapagliflozin, empagliflozin	<ul style="list-style-type: none"> <li>See labels for renal dose considerations of individual agents</li> <li>Glucose-lowering effect is lower for SGLT2 inhibitors at lower eGFR</li> </ul>	Oral	High	<ul style="list-style-type: none"> <li>DKA risk, rare in T2DM; discontinue, evaluate, and treat promptly if suspected; be aware of predisposing risk factors and clinical presentation (including euglycemic DKA); discontinue before scheduled surgery (e.g., 3–4 days), during critical illness, or during prolonged fasting to mitigate potential risk</li> <li>Increased risk of genital mycotic infections</li> <li>Necrotizing fasciitis of the perineum (Fournier gangrene), rare reports: institute prompt treatment if suspected</li> <li>Attention to volume status, blood pressure; adjust other volume-contracting agents as applicable</li> </ul>	
<b>GLP-1 RAs</b>	High to very high	No	Loss (intermediate to very high)	Benefit: dulaglutide, liraglutide, semaglutide (SQ) Neutral: exenatide once weekly, lixisenatide	Neutral	Benefit for renal endpoints in CVOTs, driven by albuminuria outcomes: dulaglutide, liraglutide, semaglutide (SQ)	<ul style="list-style-type: none"> <li>See labels for renal dose considerations of individual agents</li> <li>No dose adjustment for dulaglutide, liraglutide, semaglutide</li> <li>Monitor renal function when initiating or escalating doses in patients with renal impairment reporting severe adverse GI reactions</li> </ul>	SQ; oral (semaglutide)	High	<ul style="list-style-type: none"> <li>Risk of thyroid C-cell tumors in rodents; human relevance not determined (liraglutide, dulaglutide, exenatide extended release, semaglutide)</li> <li>Counsel patients on potential for GI side effects and their typically temporary nature; provide guidance on dietary modifications to mitigate GI side effects (reduction in meal size, mindful eating practices [e.g., stop eating once full], decreasing intake of high-fat or spicy food); consider slower dose titration for patients experiencing GI challenges</li> <li>Pancreatitis has been reported in clinical trials but causality has not been established. Discontinue if pancreatitis is suspected</li> <li>Evaluate for gallbladder disease if cholelithiasis or cholecystitis is suspected</li> </ul>	
<b>GIP and GLP-1 RA</b>	Very high	No	Loss (very high)	Under investigation	Under investigation	Under investigation	<ul style="list-style-type: none"> <li>See label for renal dose considerations</li> <li>No dose adjustment</li> <li>Monitor renal function when initiating or escalating doses in patients with renal impairment reporting severe adverse GI reactions</li> </ul>	SQ	High	<ul style="list-style-type: none"> <li>Risk of thyroid C-cell tumors in rodents; human relevance not determined</li> <li>Counsel patients on potential for GI side effects and their typically temporary nature; provide guidance on dietary modifications to mitigate GI side effects (reduction in meal size, mindful eating practices [e.g., stop eating once full], decreasing intake of high-fat or spicy food); consider slower dose titration for patients experiencing GI challenges</li> <li>Pancreatitis has been reported in clinical trials but causality has not been established. Discontinue if pancreatitis is suspected</li> <li>Evaluate for gallbladder disease if cholelithiasis or cholecystitis is suspected</li> </ul>	
<b>DPP-4 inhibitors</b>	Intermediate	No	Neutral	Neutral	Neutral (potential risk, saxagliptin)	Neutral	<ul style="list-style-type: none"> <li>Renal dose adjustment required (sitagliptin, saxagliptin, alogliptin); can be used in renal impairment</li> <li>No dose adjustment required for linagliptin</li> </ul>	Oral	High	<ul style="list-style-type: none"> <li>Pancreatitis has been reported in clinical trials but causality has not been established. Discontinue if pancreatitis is suspected</li> <li>Joint pain</li> <li>Bullous pemphigoid (postmarketing); discontinue if suspected</li> </ul>	
<b>Thiazolidinediones</b>	High	No	Gain	Potential benefit: pioglitazone	Increased risk	Neutral	<ul style="list-style-type: none"> <li>No dose adjustment required</li> <li>Generally not recommended in renal impairment due to potential for fluid retention</li> </ul>	Oral	Low	<ul style="list-style-type: none"> <li>Congestive HF (pioglitazone, rosiglitazone)</li> <li>Fluid retention (edema; heart failure)</li> <li>Benefit in NASH</li> <li>Risk of bone fractures</li> <li>Weight gain: consider lower doses to mitigate weight gain and edema</li> </ul>	
<b>Sulfonylureas (2nd generation)</b>	High	Yes	Gain	Neutral	Neutral	Neutral	<ul style="list-style-type: none"> <li>Glyburide: generally not recommended in chronic kidney disease</li> <li>Glipizide and glimepiride: initiate conservatively to avoid hypoglycemia</li> </ul>	Oral	Low	<ul style="list-style-type: none"> <li>FDA Special Warning on increased risk of CV mortality based on studies of an older sulfonylurea (tolbutamide); glimepiride shown to be CV safe (see text)</li> <li>Use with caution in persons at risk for hypoglycemia</li> </ul>	
<b>Insulin</b>	<b>Human</b>	High to very high	Yes	Gain	Neutral	Neutral	Neutral	<ul style="list-style-type: none"> <li>Lower insulin doses required with a decrease in eGFR; titrate per clinical response</li> </ul>	SQ; inhaled	Low (SQ)	<ul style="list-style-type: none"> <li>Injection site reactions</li> <li>Higher risk of hypoglycemia with human insulin (NPH or premixed formulations) vs. analogs</li> </ul>
	<b>Analog</b>								SQ	High	

CV, cardiovascular; CVOT, cardiovascular outcomes trial; DKA, diabetic ketoacidosis; DKD, diabetic kidney disease; DPP-4, dipeptidyl peptidase 4; eGFR, estimated glomerular filtration rate; FDA, U.S. Food and Drug Administration; GI, gastrointestinal; GIP, gastric inhibitory polypeptide; GLP-1 RA, glucagon-like peptide 1 receptor agonist; HF, heart failure; NASH, nonalcoholic steatohepatitis; MACE, major adverse cardiovascular events; SGLT2, sodium–glucose cotransporter 2; SQ, subcutaneous; T2DM, type 2 diabetes mellitus. \*For agent-specific dosing recommendations, please refer to manufacturers’ prescribing information. <sup>1</sup>Tsapas et al. (62). <sup>2</sup>Tsapas et al. (114). Reprinted from Davies et al. (45).